

264-1065

5/23/2013

1/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

MAY 23 2013

Ms. Laura Phelps
Bayer Cropscience
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Dear Ms. Phelps:

Subject: Fast track label amendment, add hops, signal word and KOROC statement
Ultor – Supplemental label for IR-4 new uses
EPA Registration Number 264-1065
Your submission dated 5/7/2013
OPP Decision Number D478406

The supplemental label referred to above, and submitted in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This information has been added to our files for this product.

A stamped copy of the label is enclosed for your records. Please note that this supplemental label has an expiration date of 10/15/2013. If you have any questions, contact Rita Kumar at (703) 308-8291, or kumar.rita@epa.gov.

Yours Sincerely,

A handwritten signature in black ink, appearing to read "John Hebert".

John Hebert
Product Manager 7
Insecticide-Rodenticide Branch
Registration Division (7505P)

Enclosure



Bayer CropScience

Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

ULTOR®
EPA Reg. No. 264-1065

For Use On: Hops, Citrus Fruits, Pome Fruits And Stone Fruits

This supplemental label expires on October 15th, 2013 and must not be used or distributed after this date.

Supplemental Label

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for ULTOR® attached to the container.

USE DIRECTIONS FOR SPECIFIC CROPS

HOPS

PESTS CONTROLLED	Rate fluid ounces/Acre	Rate lb ai/Acre
Hop aphid	8.0 – 10.0	0.08 – 0.10
Twospotted spider mite		
Restrictions		
Pre-Harvest Interval (PHI): 7 days.		
Minimum interval between applications: 14 days.		
Maximum ULTOR allowed per 12-month period: 20 fluid ounces/Acre (0.2 lb ai/Acre).		

**ACCEPTED
MAY 23 2013**

**Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended, for the
pesticide registered under:**

ULTOR is a Registered trademark of Bayer.
PENDING 05-06-13

EPA. Reg. No: 264-1065

CITRUS FRUITS

Crops of Crop Group 10-10 Including: Australian lime (desert, finger, and round), Brown River finger lime, Calamondin, Citron, Grapefruit, Japanese summer grapefruit, Kumquat, Lemon, Lime, Mediterranean mandarin, Mount White lime, New Guinea wild lime, Orange (sour and sweet), Pummelo, Russell River lime, Satsuma mandarin, Sweet lime, Tachibana orange, Tahiti lime, Tangelo, Tangerine, Tangor, Trifoliolate orange, Uniq fruit, including cultivars, varieties and/or hybrids of these commodities.

PESTS CONTROLLED	Rate fluid ounces/Acre	Rate lb ai/Acre
Aphids Asian citrus psyllid Broad mite California red scale Citrus leafminer Citrus bud mite Citrus red mite Citrus rust mite (silver mite) Citrus snow scale Citrus thrips Florida red scale Mealybugs Pink citrus rust mite Purple scale Texas citrus mite Whiteflies	12.0 – 16.0	0.12 – 0.16
PESTS SUPPRESSED		
Black scale Brown scale Citricola scale Cottony cushion scale Nematodes		
<p>Restrictions</p> <p>Pre-Harvest Interval (PHI): 1 day.</p> <p>Minimum interval between applications: 21 days.</p> <p>Maximum ULTOR allowed per 12-month period: 32 fluid ounces/Acre (0.32 lb ai/Acre).</p> <p>Do not apply this product within 10 days prior to bloom, during bloom, or until petal fall is complete.</p> <p>Do not apply non-ionic surfactants in tank mix combination with ULTOR on white grapefruit.</p>		

POME FRUITS

Crops of Crop Group 11-10 Including: Apple, Asian pear, Azarole, Chinese quince, Crabapple, Japanese quince, Loquat, Mayhaw, Medlar, Pear, Quince, Tejocote, including cultivars, varieties and/or hybrids of these commodities.

PESTS CONTROLLED	Rate fluid ounces/Acre	Rate lb ai/Acre
Aphids (including Woolly Apple Aphid) Apple rust mite Mealybugs Pear psylla Pear rust mite San Jose scale Whiteflies	8.0 – 14.0	0.08 – 0.14
PESTS SUPPRESSED		
Apple gall midge Codling moth European red mite Micro-lepidoptera leafminers Pear leaf midge Twospotted spider mite White apple leafhopper		
<p>Restrictions</p> <p>Pre-Harvest Interval (PHI): 7 days.</p> <p>Minimum interval between applications: 14 days.</p> <p>Maximum ULTOR allowed per 12 month period: 40 fluid ounces/Acre (0.4 lb ai/Acre).</p> <p>Do not apply until after petal fall.</p> <p>For control of San Jose scale west of the Rocky Mountains [including all of MT, WY, CO, and NM],, apply immediately after petal fall, followed by a second application 14 – 21 days later. For control of San Jose scale east of the Rocky Mountains, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.</p>		

STONE FRUITS

Crops of Crop Group 12 Including: Apricot, Cherry (sweet and tart), Nectarine, Peach, Plum (includes Chickasaw, Damson, and Japanese), Plumcot, Prune (fresh and dried)

PESTS CONTROLLED	Rate fluid ounces/Acre	Rate lb ai/Acre
Aphids Mealybugs San Jose scale White peach scale Whiteflies		
PESTS SUPPRESSED	8.0 – 14.0	0.08 – 0.14
Black scale Cherry fruit fly European fruit lecanium scale European red mite Nematodes Twospotted spider mite		
<p>Restrictions</p> <p>Pre-Harvest Interval (PHI): 7 days.</p> <p>Minimum interval between applications: 14 days.</p> <p>Maximum ULTOR allowed per 12-month period: 24 fluid ounces/Acre (0.24 lb ai/Acre).</p> <p>Do not apply until after petal fall.</p> <p>For control of San Jose scale, apply immediately after petal fall; under heavy infestation pressure or where difficult control conditions exist, a second application may be necessary.</p>		

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).